




## 3.10 Exercises

1–4 Find the linearization  $L(x)$  of the function at  $a$ .

1.  $f(x) = x^4 + 3x^2$ ,  $a = -1$       2.  $f(x) = \sin x$ ,  $a = \pi/6$   
 3.  $f(x) = \sqrt{x}$ ,  $a = 4$       4.  $f(x) = x^{3/4}$ ,  $a = 16$

 5. Find the linear approximation of the function  $f(x) = \sqrt{1-x}$  at  $a = 0$  and use it to approximate the numbers  $\sqrt{0.9}$  and  $\sqrt{0.99}$ . Illustrate by graphing  $f$  and the tangent line.

 6. Find the linear approximation of the function  $g(x) = \sqrt[3]{1+x}$  at  $a = 0$  and use it to approximate the numbers  $\sqrt[3]{0.95}$  and  $\sqrt[3]{1.1}$ . Illustrate by graphing  $g$  and the tangent line.

 7–10 Verify the given linear approximation at  $a = 0$ . Then determine the values of  $x$  for which the linear approximation is accurate to within 0.1.

7.  $\ln(1+x) \approx x$       8.  $(1+x)^{-3} \approx 1-3x$   
 9.  $\sqrt[4]{1+2x} \approx 1 + \frac{1}{2}x$       10.  $e^x \cos x \approx 1+x$

11–14 Find the differential of each function.

11. (a)  $y = x^2 \sin 2x$       (b)  $y = \ln \sqrt{1+t^2}$   
 12. (a)  $y = s/(1+2s)$       (b)  $y = e^{-u} \cos u$   
 13. (a)  $y = \tan \sqrt{t}$       (b)  $y = \frac{1-v^2}{1+v^2}$   
 14. (a)  $y = e^{\tan \pi t}$       (b)  $y = \sqrt{1 + \ln z}$

15–18 (a) Find the differential  $dy$  and (b) evaluate  $dy$  for the given values of  $x$  and  $dx$ .

15.  $y = e^{x/10}$ ,  $x = 0$ ,  $dx = 0.1$   
 16.  $y = \cos \pi x$ ,  $x = \frac{1}{3}$ ,  $dx = -0.02$   
 17.  $y = \sqrt{3+x^2}$ ,  $x = 1$ ,  $dx = -0.1$   
 18.  $y = \frac{x+1}{x-1}$ ,  $x = 2$ ,  $dx = 0.05$

19–22 Compute  $\Delta y$  and  $dy$  for the given values of  $x$  and  $dx = \Delta x$ . Then sketch a diagram like Figure 5 showing the line segments with lengths  $dx$ ,  $dy$ , and  $\Delta y$ .

19.  $y = 2x - x^2$ ,  $x = 2$ ,  $\Delta x = -0.4$   
 20.  $y = \sqrt{x}$ ,  $x = 1$ ,  $\Delta x = 1$   
 21.  $y = 2/x$ ,  $x = 4$ ,  $\Delta x = 1$   
 22.  $y = e^x$ ,  $x = 0$ ,  $\Delta x = 0.5$

23–28 Use a linear approximation (or differentials) to estimate the given number.

23.  $(1.999)^4$       24.  $e^{-0.015}$

25.  $\sqrt[3]{1001}$       26.  $1/4.002$   
 27.  $\tan 44^\circ$       28.  $\sqrt{99.8}$

29–31 Explain, in terms of linear approximations or differentials, why the approximation is reasonable.

29.  $\sec 0.08 \approx 1$       30.  $(1.01)^6 \approx 1.06$   
 31.  $\ln 1.05 \approx 0.05$

32. Let  $f(x) = (x-1)^2$        $g(x) = e^{-2x}$   
 and  $h(x) = 1 + \ln(1-2x)$

- (a) Find the linearizations of  $f$ ,  $g$ , and  $h$  at  $a = 0$ . What do you notice? How do you explain what happened?  
 (b) Graph  $f$ ,  $g$ , and  $h$  and their linear approximations. For which function is the linear approximation best? For which is it worst? Explain.



33. The edge of a cube was found to be 30 cm with a possible error in measurement of 0.1 cm. Use differentials to estimate the maximum possible error, relative error, and percentage error in computing (a) the volume of the cube and (b) the surface area of the cube.
34. The radius of a circular disk is given as 24 cm with a maximum error in measurement of 0.2 cm.  
 (a) Use differentials to estimate the maximum error in the calculated area of the disk.  
 (b) What is the relative error? What is the percentage error?
35. The circumference of a sphere was measured to be 84 cm with a possible error of 0.5 cm.  
 (a) Use differentials to estimate the maximum error in the calculated surface area. What is the relative error?  
 (b) Use differentials to estimate the maximum error in the calculated volume. What is the relative error?
36. Use differentials to estimate the amount of paint needed to apply a coat of paint 0.05 cm thick to a hemispherical dome with diameter 50 m.
37. (a) Use differentials to find a formula for the approximate volume of a thin cylindrical shell with height  $h$ , inner radius  $r$ , and thickness  $\Delta r$ .  
 (b) What is the error involved in using the formula from part (a)?
38. One side of a right triangle is known to be 20 cm long and the opposite angle is measured as  $30^\circ$ , with a possible error of  $\pm 1^\circ$ .  
 (a) Use differentials to estimate the error in computing the length of the hypotenuse.  
 (b) What is the percentage error?

39. If a current  $I$  passes through a resistor with resistance  $R$ , Ohm's Law states that the voltage drop is  $V = RI$ . If  $V$  is constant and  $R$  is measured with a certain error, use differentials to show that the relative error in calculating  $I$  is approximately the same (in magnitude) as the relative error in  $R$ .
40. When blood flows along a blood vessel, the flux  $F$  (the volume of blood per unit time that flows past a given point) is proportional to the fourth power of the radius  $R$  of the blood vessel:

$$F = kR^4$$

(This is known as Poiseuille's Law; we will show why it is true in Section 8.4.) A partially clogged artery can be expanded by an operation called angioplasty, in which a balloon-tipped catheter is inflated inside the artery in order to widen it and restore the normal blood flow.

Show that the relative change in  $F$  is about four times the relative change in  $R$ . How will a 5% increase in the radius affect the flow of blood?

41. Establish the following rules for working with differentials (where  $c$  denotes a constant and  $u$  and  $v$  are functions of  $x$ ).

$$\begin{array}{ll} \text{(a) } dc = 0 & \text{(b) } d(cu) = c \, du \\ \text{(c) } d(u + v) = du + dv & \text{(d) } d(uv) = u \, dv + v \, du \\ \text{(e) } d\left(\frac{u}{v}\right) = \frac{v \, du - u \, dv}{v^2} & \text{(f) } d(x^n) = nx^{n-1} \, dx \end{array}$$

42. On page 431 of *Physics: Calculus*, 2d ed., by Eugene Hecht (Pacific Grove, CA, 2000), in the course of deriving the formula  $T = 2\pi\sqrt{L/g}$  for the period of a pendulum of length  $L$ , the author obtains the equation  $a_T = -g \sin \theta$  for the tangen-

tial acceleration of the bob of the pendulum. He then says, "for small angles, the value of  $\theta$  in radians is very nearly the value of  $\sin \theta$ ; they differ by less than 2% out to about  $20^\circ$ ."


- (a) Verify the linear approximation at 0 for the sine function:

$$\sin x \approx x$$

- (b) Use a graphing device to determine the values of  $x$  for which  $\sin x$  and  $x$  differ by less than 2%. Then verify Hecht's statement by converting from radians to degrees.

43. Suppose that the only information we have about a function  $f$  is that  $f(1) = 5$  and the graph of its derivative is as shown.

- (a) Use a linear approximation to estimate  $f(0.9)$  and  $f(1.1)$ .  
 (b) Are your estimates in part (a) too large or too small? Explain.




44. Suppose that we don't have a formula for  $g(x)$  but we know that  $g(2) = -4$  and  $g'(x) = \sqrt{x^2 + 5}$  for all  $x$ .  
 (a) Use a linear approximation to estimate  $g(1.95)$  and  $g(2.05)$ .  
 (b) Are your estimates in part (a) too large or too small? Explain.

## LABORATORY PROJECT TAYLOR POLYNOMIALS

The tangent line approximation  $L(x)$  is the best first-degree (linear) approximation to  $f(x)$  near  $x = a$  because  $f(x)$  and  $L(x)$  have the same rate of change (derivative) at  $a$ . For a better approximation than a linear one, let's try a second-degree (quadratic) approximation  $P(x)$ . In other words, we approximate a curve by a parabola instead of by a straight line. To make sure that the approximation is a good one, we stipulate the following:

- (i)  $P(a) = f(a)$  ( $P$  and  $f$  should have the same value at  $a$ .)
- (ii)  $P'(a) = f'(a)$  ( $P$  and  $f$  should have the same rate of change at  $a$ .)
- (iii)  $P''(a) = f''(a)$  (The slopes of  $P$  and  $f$  should change at the same rate at  $a$ .)

1. Find the quadratic approximation  $P(x) = A + Bx + Cx^2$  to the function  $f(x) = \cos x$  that satisfies conditions (i), (ii), and (iii) with  $a = 0$ . Graph  $P$ ,  $f$ , and the linear approximation  $L(x) = 1$  on a common screen. Comment on how well the functions  $P$  and  $L$  approximate  $f$ .
2. Determine the values of  $x$  for which the quadratic approximation  $f(x) \approx P(x)$  in Problem 1 is accurate to within 0.1. [Hint: Graph  $y = P(x)$ ,  $y = \cos x - 0.1$ , and  $y = \cos x + 0.1$  on a common screen.]

 Graphing calculator or computer required