

**SOLUTION 2** From Equation 3 (proved in Example 3), we have

$$\begin{aligned} \frac{d}{dx}(\sinh^{-1}x) &= \frac{d}{dx} \ln(x + \sqrt{x^2 + 1}) \\ &= \frac{1}{x + \sqrt{x^2 + 1}} \frac{d}{dx}(x + \sqrt{x^2 + 1}) \\ &= \frac{1}{x + \sqrt{x^2 + 1}} \left( 1 + \frac{x}{\sqrt{x^2 + 1}} \right) \\ &= \frac{\sqrt{x^2 + 1} + x}{(x + \sqrt{x^2 + 1})\sqrt{x^2 + 1}} \\ &= \frac{1}{\sqrt{x^2 + 1}} \end{aligned}$$

**V EXAMPLE 5** Find  $\frac{d}{dx}[\tanh^{-1}(\sin x)]$ .

**SOLUTION** Using Table 6 and the Chain Rule, we have

$$\begin{aligned} \frac{d}{dx}[\tanh^{-1}(\sin x)] &= \frac{1}{1 - (\sin x)^2} \frac{d}{dx}(\sin x) \\ &= \frac{1}{1 - \sin^2 x} \cos x = \frac{\cos x}{\cos^2 x} = \sec x \end{aligned}$$

### 3.11 Exercises

**1–6** Find the numerical value of each expression.

- |                                |                    |
|--------------------------------|--------------------|
| 1. (a) $\sinh 0$               | (b) $\cosh 0$      |
| 2. (a) $\tanh 0$               | (b) $\tanh 1$      |
| 3. (a) $\sinh(\ln 2)$          | (b) $\sinh 2$      |
| 4. (a) $\cosh 3$               | (b) $\cosh(\ln 3)$ |
| 5. (a) $\operatorname{sech} 0$ | (b) $\cosh^{-1} 1$ |
| 6. (a) $\sinh 1$               | (b) $\sinh^{-1} 1$ |

**7–19** Prove the identity.

- $\sinh(-x) = -\sinh x$   
(This shows that  $\sinh$  is an odd function.)
- $\cosh(-x) = \cosh x$   
(This shows that  $\cosh$  is an even function.)
- $\cosh x + \sinh x = e^x$
- $\cosh x - \sinh x = e^{-x}$
- $\sinh(x + y) = \sinh x \cosh y + \cosh x \sinh y$
- $\cosh(x + y) = \cosh x \cosh y + \sinh x \sinh y$
- $\coth^2 x - 1 = \operatorname{csch}^2 x$
- $\tanh(x + y) = \frac{\tanh x + \tanh y}{1 + \tanh x \tanh y}$
- $\sinh 2x = 2 \sinh x \cosh x$
- $\cosh 2x = \cosh^2 x + \sinh^2 x$
- $\tanh(\ln x) = \frac{x^2 - 1}{x^2 + 1}$
- $\frac{1 + \tanh x}{1 - \tanh x} = e^{2x}$
- $(\cosh x + \sinh x)^n = \cosh nx + \sinh nx$   
( $n$  any real number)
- If  $\tanh x = \frac{12}{13}$ , find the values of the other hyperbolic functions at  $x$ .
- If  $\cosh x = \frac{5}{3}$  and  $x > 0$ , find the values of the other hyperbolic functions at  $x$ .
- (a) Use the graphs of  $\sinh$ ,  $\cosh$ , and  $\tanh$  in Figures 1–3 to draw the graphs of  $\operatorname{csch}$ ,  $\operatorname{sech}$ , and  $\coth$ .





- (b) Check the graphs that you sketched in part (a) by using a graphing device to produce them.

23. Use the definitions of the hyperbolic functions to find each of the following limits.

- |                                                          |                                                         |
|----------------------------------------------------------|---------------------------------------------------------|
| (a) $\lim_{x \rightarrow \infty} \tanh x$                | (b) $\lim_{x \rightarrow -\infty} \tanh x$              |
| (c) $\lim_{x \rightarrow \infty} \sinh x$                | (d) $\lim_{x \rightarrow -\infty} \sinh x$              |
| (e) $\lim_{x \rightarrow \infty} \operatorname{sech} x$  | (f) $\lim_{x \rightarrow \infty} \operatorname{coth} x$ |
| (g) $\lim_{x \rightarrow 0^+} \operatorname{coth} x$     | (h) $\lim_{x \rightarrow 0^-} \operatorname{coth} x$    |
| (i) $\lim_{x \rightarrow -\infty} \operatorname{csch} x$ |                                                         |

24. Prove the formulas given in Table 1 for the derivatives of the functions (a)  $\cosh$ , (b)  $\tanh$ , (c)  $\operatorname{csch}$ , (d)  $\operatorname{sech}$ , and (e)  $\operatorname{coth}$ .

25. Give an alternative solution to Example 3 by letting  $y = \sinh^{-1}x$  and then using Exercise 9 and Example 1(a) with  $x$  replaced by  $y$ .

26. Prove Equation 4.

27. Prove Equation 5 using (a) the method of Example 3 and (b) Exercise 18 with  $x$  replaced by  $y$ .

28. For each of the following functions (i) give a definition like those in [\[2\]](#), (ii) sketch the graph, and (iii) find a formula similar to Equation 3.

- (a)  $\operatorname{csch}^{-1}$       (b)  $\operatorname{sech}^{-1}$       (c)  $\operatorname{coth}^{-1}$

29. Prove the formulas given in Table 6 for the derivatives of the following functions.

- (a)  $\cosh^{-1}$       (b)  $\tanh^{-1}$       (c)  $\operatorname{csch}^{-1}$   
 (d)  $\operatorname{sech}^{-1}$       (e)  $\operatorname{coth}^{-1}$

30–45 Find the derivative. Simplify where possible.

- |                                                                   |                                              |
|-------------------------------------------------------------------|----------------------------------------------|
| 30. $f(x) = \tanh(1 + e^{2x})$                                    | 31. $f(x) = x \sinh x - \cosh x$             |
| 32. $g(x) = \cosh(\ln x)$                                         | 33. $h(x) = \ln(\cosh x)$                    |
| 34. $y = x \operatorname{coth}(1 + x^2)$                          | 35. $y = e^{\cosh 3x}$                       |
| 36. $f(t) = \operatorname{csch} t(1 - \ln \operatorname{csch} t)$ | 37. $f(t) = \operatorname{sech}^2(e^t)$      |
| 38. $y = \sinh(\cosh x)$                                          | 39. $G(x) = \frac{1 - \cosh x}{1 + \cosh x}$ |
| 40. $y = \sinh^{-1}(\tan x)$                                      | 41. $y = \cosh^{-1}\sqrt{x}$                 |
| 42. $y = x \tanh^{-1}x + \ln \sqrt{1 - x^2}$                      |                                              |
| 43. $y = x \sinh^{-1}(x/3) - \sqrt{9 + x^2}$                      |                                              |
| 44. $y = \operatorname{sech}^{-1}(e^{-x})$                        |                                              |
| 45. $y = \operatorname{coth}^{-1}(\sec x)$                        |                                              |

46. Show that  $\frac{d}{dx} \sqrt[4]{\frac{1 + \tanh x}{1 - \tanh x}} = \frac{1}{2}e^{x/2}$

47. Show that  $\frac{d}{dx} \arctan(\tanh x) = \operatorname{sech} 2x$ .

48. The Gateway Arch in St. Louis was designed by Eero Saarinen and was constructed using the equation

$$y = 211.49 - 20.96 \cosh 0.03291765x$$

for the central curve of the arch, where  $x$  and  $y$  are measured in meters and  $|x| \leq 91.20$ .



- (a) Graph the central curve.  
 (b) What is the height of the arch at its center?  
 (c) At what points is the height 100 m?  
 (d) What is the slope of the arch at the points in part (c)?

49. If a water wave with length  $L$  moves with velocity  $v$  in a body of water with depth  $d$ , then

$$v = \sqrt{\frac{gL}{2\pi} \tanh\left(\frac{2\pi d}{L}\right)}$$

where  $g$  is the acceleration due to gravity. (See Figure 5.) Explain why the approximation

$$v \approx \sqrt{\frac{gL}{2\pi}}$$


is appropriate in deep water.



50. A flexible cable always hangs in the shape of a catenary  $y = c + a \cosh(x/a)$ , where  $c$  and  $a$  are constants and  $a > 0$  (see Figure 4 and Exercise 52). Graph several members of the family of functions  $y = a \cosh(x/a)$ . How does the graph change as  $a$  varies?

51. A telephone line hangs between two poles 14 m apart in the shape of the catenary  $y = 20 \cosh(x/20) - 15$ , where  $x$  and  $y$  are measured in meters.

- (a) Find the slope of this curve where it meets the right pole.  
 (b) Find the angle  $\theta$  between the line and the pole.



52. Using principles from physics it can be shown that when a cable is hung between two poles, it takes the shape of a curve  $y = f(x)$  that satisfies the differential equation

$$\frac{d^2y}{dx^2} = \frac{\rho g}{T} \sqrt{1 + \left(\frac{dy}{dx}\right)^2}$$

where  $\rho$  is the linear density of the cable,  $g$  is the acceleration due to gravity,  $T$  is the tension in the cable at its lowest point, and the coordinate system is chosen appropriately. Verify that the function

$$y = f(x) = \frac{T}{\rho g} \cosh\left(\frac{\rho g x}{T}\right)$$

is a solution of this differential equation.

53. A cable with linear density  $\rho = 2$  kg/m is strung from the tops of two poles that are 200 m apart.
- (a) Use Exercise 52 to find the tension  $T$  so that the cable is 60 m above the ground at its lowest point. How tall are the poles?
- (b) If the tension is doubled, what is the new low point of the cable? How tall are the poles now?

54. Evaluate  $\lim_{x \rightarrow \infty} \frac{\sinh x}{e^x}$ .

55. (a) Show that any function of the form


$$y = A \sinh mx + B \cosh mx$$

satisfies the differential equation  $y'' = m^2 y$ .

- (b) Find  $y = y(x)$  such that  $y'' = 9y$ ,  $y(0) = -4$ , and  $y'(0) = 6$ .

56. If  $x = \ln(\sec \theta + \tan \theta)$ , show that  $\sec \theta = \cosh x$ .

57. At what point of the curve  $y = \cosh x$  does the tangent have slope 1?

-  58. Investigate the family of functions

$$f_n(x) = \tanh(n \sin x)$$

where  $n$  is a positive integer. Describe what happens to the graph of  $f_n$  when  $n$  becomes large.

59. Show that if  $a \neq 0$  and  $b \neq 0$ , then there exist numbers  $\alpha$  and  $\beta$  such that  $ae^x + be^{-x}$  equals either  $\alpha \sinh(x + \beta)$  or  $\alpha \cosh(x + \beta)$ . In other words, almost every function of the form  $f(x) = ae^x + be^{-x}$  is a shifted and stretched hyperbolic sine or cosine function.

### 3 Review

#### Concept Check

- State each differentiation rule both in symbols and in words.
  - The Power Rule
  - The Constant Multiple Rule
  - The Sum Rule
  - The Difference Rule
  - The Product Rule
  - The Quotient Rule
  - The Chain Rule
- State the derivative of each function.
 

(a) $y = x^n$	(b) $y = e^x$	(c) $y = a^x$
(d) $y = \ln x$	(e) $y = \log_a x$	(f) $y = \sin x$
(g) $y = \cos x$	(h) $y = \tan x$	(i) $y = \csc x$
(j) $y = \sec x$	(k) $y = \cot x$	(l) $y = \sin^{-1} x$
(m) $y = \cos^{-1} x$	(n) $y = \tan^{-1} x$	(o) $y = \sinh x$
(p) $y = \cosh x$	(q) $y = \tanh x$	(r) $y = \sinh^{-1} x$
(s) $y = \cosh^{-1} x$	(t) $y = \tanh^{-1} x$	
- How is the number  $e$  defined?
  - Express  $e$  as a limit.
  - Why is the natural exponential function  $y = e^x$  used more often in calculus than the other exponential functions  $y = a^x$ ?
- Why is the natural logarithmic function  $y = \ln x$  used more often in calculus than the other logarithmic functions  $y = \log_a x$ ?
- Explain how implicit differentiation works.
  - Explain how logarithmic differentiation works.
- Give several examples of how the derivative can be interpreted as a rate of change in physics, chemistry, biology, economics, or other sciences.
- Write a differential equation that expresses the law of natural growth.
  - Under what circumstances is this an appropriate model for population growth?
  - What are the solutions of this equation?
- Write an expression for the linearization of  $f$  at  $a$ .
  - If  $y = f(x)$ , write an expression for the differential  $dy$ .
  - If  $dx = \Delta x$ , draw a picture showing the geometric meanings of  $\Delta y$  and  $dy$ .

#### True-False Quiz

Determine whether the statement is true or false. If it is true, explain why. If it is false, explain why or give an example that disproves the statement.

1. If  $f$  and  $g$  are differentiable, then

$$\frac{d}{dx} [f(x) + g(x)] = f'(x) + g'(x)$$

2. If  $f$  and  $g$  are differentiable, then

$$\frac{d}{dx} [f(x)g(x)] = f'(x)g'(x)$$

3. If  $f$  and  $g$  are differentiable, then

$$\frac{d}{dx} [f(g(x))] = f'(g(x))g'(x)$$

4. If  $f$  is differentiable, then  $\frac{d}{dx} \sqrt{f(x)} = \frac{f'(x)}{2\sqrt{f(x)}}$ .

5. If  $f$  is differentiable, then  $\frac{d}{dx} f(\sqrt{x}) = \frac{f'(x)}{2\sqrt{x}}$ .

6. If  $y = e^2$ , then  $y' = 2e$ .