

$$\text{so } \frac{\Delta y}{\Delta x} = [f'(b) + \varepsilon_2][g'(a) + \varepsilon_1]$$

As  $\Delta x \rightarrow 0$ , Equation 8 shows that  $\Delta u \rightarrow 0$ . So both  $\varepsilon_1 \rightarrow 0$  and  $\varepsilon_2 \rightarrow 0$  as  $\Delta x \rightarrow 0$ . Therefore

$$\begin{aligned} \frac{dy}{dx} &= \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x} = \lim_{\Delta x \rightarrow 0} [f'(b) + \varepsilon_2][g'(a) + \varepsilon_1] \\ &= f'(b)g'(a) = f'(g(a))g'(a) \end{aligned}$$

This proves the Chain Rule.

### 3.4 Exercises

**1–6** Write the composite function in the form  $f(g(x))$ . [Identify the inner function  $u = g(x)$  and the outer function  $y = f(u)$ .] Then find the derivative  $dy/dx$ .

- |                           |                         |
|---------------------------|-------------------------|
| 1. $y = \sqrt[3]{1 + 4x}$ | 2. $y = (2x^3 + 5)^4$   |
| 3. $y = \tan \pi x$       | 4. $y = \sin(\cot x)$   |
| 5. $y = e^{\sqrt{x}}$     | 6. $y = \sqrt{2 - e^x}$ |

**7–46** Find the derivative of the function.

- |  |                                       |
|--|---------------------------------------|
| 7. $F(x) = (x^4 + 3x^2 - 2)^5$         | 8. $F(x) = (4x - x^2)^{100}$          |
| 9. $F(x) = \sqrt{1 - 2x}$              | 10. $f(x) = \frac{1}{(1 + \sec x)^2}$ |
| 11. $f(z) = \frac{1}{z^2 + 1}$         | 12. $f(t) = \sin(e^t) + e^{\sin t}$   |
| 13. $y = \cos(a^3 + x^3)$              | 14. $y = a^3 + \cos^3 x$              |
| 15. $y = xe^{-kx}$                     | 16. $y = e^{-2t} \cos 4t$             |
| 17. $f(x) = (2x - 3)^4(x^2 + x + 1)^5$ |                                       |
| 18. $g(x) = (x^2 + 1)^3(x^2 + 2)^6$    |                                       |
| 19. $h(t) = (t + 1)^{2/3}(2t^2 - 1)^3$ |                                       |
| 20. $F(t) = (3t - 1)^4(2t + 1)^{-3}$   |                                       |

21. $y = \left(\frac{x^2 + 1}{x^2 - 1}\right)^3$	22. $f(s) = \sqrt{\frac{s^2 + 1}{s^2 + 4}}$
--	---

23. $y = \sqrt{1 + 2e^{3x}}$	24. $y = 10^{1-x^2}$
------------------------------	----------------------

25. $y = 5^{-1/x}$	26. $G(y) = \frac{(y - 1)^4}{(y^2 + 2y)^5}$
--------------------	---

27. $y = \frac{r}{\sqrt{r^2 + 1}}$	28. $y = \frac{e^u - e^{-u}}{e^u + e^{-u}}$
------------------------------------	---

29. $F(t) = e^{t \sin 2t}$	30. $F(v) = \left(\frac{v}{v^3 + 1}\right)^6$
----------------------------	---

31. $y = \sin(\tan 2x)$	32. $y = \sec^2(m\theta)$
-------------------------	---------------------------

33. $y = 2^{\sin \pi x}$	34. $y = x^2 e^{-1/x}$
--------------------------	------------------------

35. $y = \cos\left(\frac{1 - e^{2x}}{1 + e^{2x}}\right)$	36. $y = \sqrt{1 + xe^{-2x}}$
--	-------------------------------

37. $y = \cot^2(\sin \theta)$	38. $y = e^{k \tan \sqrt{x}}$
-------------------------------	-------------------------------

39. $f(t) = \tan(e^t) + e^{\tan t}$	40. $y = \sin(\sin(\sin x))$
-------------------------------------	------------------------------

41. $f(t) = \sin^2(e^{\sin^2 t})$	42. $y = \sqrt{x + \sqrt{x + \sqrt{x}}}$
-----------------------------------	--

43. $g(x) = (2ra^{rx} + n)^p$	44. $y = 2^{3x^2}$
-------------------------------	--------------------

45. $y = \cos \sqrt{\sin(\tan \pi x)}$	46. $y = [x + (x + \sin^2 x)^3]^4$
--	------------------------------------

**47–50** Find  $y'$  and  $y''$ .

47. $y = \cos(x^2)$	48. $y = \cos^2 x$
---------------------	--------------------


49. $y = e^{\alpha x} \sin \beta x$	50. $y = e^{e^x}$
-------------------------------------	-------------------

**51–54** Find an equation of the tangent line to the curve at the given point.


51. $y = (1 + 2x)^{10}, (0, 1)$	52. $y = \sqrt{1 + x^3}, (2, 3)$
---------------------------------	----------------------------------

53. $y = \sin(\sin x), (\pi, 0)$	54. $y = \sin x + \sin^2 x, (0, 0)$
----------------------------------	-------------------------------------


**55.** (a) Find an equation of the tangent line to the curve  $y = 2/(1 + e^{-x})$  at the point  $(0, 1)$ .

 (b) Illustrate part (a) by graphing the curve and the tangent line on the same screen.

**56.** (a) The curve  $y = |x|/\sqrt{2 - x^2}$  is called a *bullet-nose curve*. Find an equation of the tangent line to this curve at the point  $(1, 1)$ .

 (b) Illustrate part (a) by graphing the curve and the tangent line on the same screen.

**57.** (a) If  $f(x) = x\sqrt{2 - x^2}$ , find  $f'(x)$ .

 (b) Check to see that your answer to part (a) is reasonable by comparing the graphs of  $f$  and  $f'$ .

58. The function  $f(x) = \sin(x + \sin 2x)$ ,  $0 \leq x \leq \pi$ , arises in applications to frequency modulation (FM) synthesis.
- Use a graph of  $f$  produced by a graphing device to make a rough sketch of the graph of  $f'$ .
  - Calculate  $f'(x)$  and use this expression, with a graphing device, to graph  $f'$ . Compare with your sketch in part (a).

59. Find all points on the graph of the function  $f(x) = 2 \sin x + \sin^2 x$  at which the tangent line is horizontal.

60. Find the  $x$ -coordinates of all points on the curve  $y = \sin 2x - 2 \sin x$  at which the tangent line is horizontal.

61. If  $F(x) = f(g(x))$ , where  $f(-2) = 8$ ,  $f'(-2) = 4$ ,  $f'(5) = 3$ ,  $g(5) = -2$ , and  $g'(5) = 6$ , find  $F'(5)$ .

62. If  $h(x) = \sqrt{4 + 3f(x)}$ , where  $f(1) = 7$  and  $f'(1) = 4$ , find  $h'(1)$ .

63. A table of values for  $f$ ,  $g$ ,  $f'$ , and  $g'$  is given.


$x$	$f(x)$	$g(x)$	$f'(x)$	$g'(x)$
1	3	2	4	6
2	1	8	5	7
3	7	2	7	9

- If  $h(x) = f(g(x))$ , find  $h'(1)$ .
- If  $H(x) = g(f(x))$ , find  $H'(1)$ .


64. Let  $f$  and  $g$  be the functions in Exercise 63.

- If  $F(x) = f(f(x))$ , find  $F'(2)$ .
- If  $G(x) = g(g(x))$ , find  $G'(3)$ .


65. If  $f$  and  $g$  are the functions whose graphs are shown, let  $u(x) = f(g(x))$ ,  $v(x) = g(f(x))$ , and  $w(x) = g(g(x))$ . Find each derivative, if it exists. If it does not exist, explain why.
- $u'(1)$
  - $v'(1)$
  - $w'(1)$



66. If  $f$  is the function whose graph is shown, let  $h(x) = f(f(x))$  and  $g(x) = f(x^2)$ . Use the graph of  $f$  to estimate the value of each derivative.
- $h'(2)$
  - $g'(2)$



67. If  $g(x) = \sqrt{f(x)}$ , where the graph of  $f$  is shown, evaluate  $g'(3)$ .



68. Suppose  $f$  is differentiable on  $\mathbb{R}$  and  $\alpha$  is a real number. Let  $F(x) = f(x^\alpha)$  and  $G(x) = [f(x)]^\alpha$ . Find expressions for (a)  $F'(x)$  and (b)  $G'(x)$ .

69. Suppose  $f$  is differentiable on  $\mathbb{R}$ . Let  $F(x) = f(e^x)$  and  $G(x) = e^{f(x)}$ . Find expressions for (a)  $F'(x)$  and (b)  $G'(x)$ .

70. Let  $g(x) = e^{kx} + f(x)$  and  $h(x) = e^{kx}f(x)$ , where  $f(0) = 3$ ,  $f'(0) = 5$ , and  $f''(0) = -2$ .

- Find  $g'(0)$  and  $g''(0)$  in terms of  $c$ .
- In terms of  $k$ , find an equation of the tangent line to the graph of  $h$  at the point where  $x = 0$ .

71. Let  $r(x) = f(g(h(x)))$ , where  $h(1) = 2$ ,  $g(2) = 3$ ,  $h'(1) = 4$ ,  $g'(2) = 5$ , and  $f'(3) = 6$ . Find  $r'(1)$ .

72. If  $g$  is a twice differentiable function and  $f(x) = xg(x^2)$ , find  $f''$  in terms of  $g$ ,  $g'$ , and  $g''$ .

73. If  $F(x) = f(3f(4f(x)))$ , where  $f(0) = 0$  and  $f'(0) = 2$ , find  $F'(0)$ .

74. If  $F(x) = f(xf(xf(x)))$ , where  $f(1) = 2$ ,  $f(2) = 3$ ,  $f'(1) = 4$ ,  $f'(2) = 5$ , and  $f'(3) = 6$ , find  $F'(1)$ .

75. Show that the function  $y = e^{2x}(A \cos 3x + B \sin 3x)$  satisfies the differential equation  $y'' - 4y' + 13y = 0$ .

76. For what values of  $r$  does the function  $y = e^{rx}$  satisfy the differential equation  $y'' - 4y' + y = 0$ ?

77. Find the 50th derivative of  $y = \cos 2x$ .

78. Find the 1000th derivative of  $f(x) = xe^{-x}$ .

79. The displacement of a particle on a vibrating string is given by the equation  $s(t) = 10 + \frac{1}{4} \sin(10\pi t)$  where  $s$  is measured in centimeters and  $t$  in seconds. Find the velocity of the particle after  $t$  seconds.

80. If the equation of motion of a particle is given by  $s = A \cos(\omega t + \delta)$ , the particle is said to undergo *simple harmonic motion*.

- Find the velocity of the particle at time  $t$ .
- When is the velocity 0?

81. A Cepheid variable star is a star whose brightness alternately increases and decreases. The most easily visible such star is Delta Cephei, for which the interval between times of maximum brightness is 5.4 days. The average brightness of this star is 4.0 and its brightness changes by  $\pm 0.35$ . In view of these data, the brightness of Delta Cephei at time  $t$ , where  $t$  is mea-

sured in days, has been modeled by the function


$$B(t) = 4.0 + 0.35 \sin\left(\frac{2\pi t}{5.4}\right)$$

- (a) Find the rate of change of the brightness after  $t$  days.  
 (b) Find, correct to two decimal places, the rate of increase after one day.

- 82.** In Example 4 in Section 1.3 we arrived at a model for the length of daylight (in hours) in Philadelphia on the  $t$ th day of the year:

$$L(t) = 12 + 2.8 \sin\left[\frac{2\pi}{365}(t - 80)\right]$$

Use this model to compare how the number of hours of daylight is increasing in Philadelphia on March 21 and May 21.

-  **83.** The motion of a spring that is subject to a frictional force or a damping force (such as a shock absorber in a car) is often modeled by the product of an exponential function and a sine or cosine function. Suppose the equation of motion of a point on such a spring is


$$s(t) = 2e^{-1.5t} \sin 2\pi t$$

where  $s$  is measured in centimeters and  $t$  in seconds. Find the velocity after  $t$  seconds and graph both the position and velocity functions for  $0 \leq t \leq 2$ .

- 84.** Under certain circumstances a rumor spreads according to the equation

$$p(t) = \frac{1}{1 + ae^{-kt}}$$

where  $p(t)$  is the proportion of the population that knows the rumor at time  $t$  and  $a$  and  $k$  are positive constants. [In Section 9.4 we will see that this is a reasonable equation for  $p(t)$ .]


- (a) Find  $\lim_{t \rightarrow \infty} p(t)$ .  
 (b) Find the rate of spread of the rumor.  
 (c) Graph  $p$  for the case  $a = 10$ ,  $k = 0.5$  with  $t$  measured in hours. Use the graph to estimate how long it will take for 80% of the population to hear the rumor.

- 85.** A particle moves along a straight line with displacement  $s(t)$ , velocity  $v(t)$ , and acceleration  $a(t)$ . Show that

$$a(t) = v(t) \frac{dv}{ds}$$

Explain the difference between the meanings of the derivatives  $dv/dt$  and  $dv/ds$ .

- 86.** Air is being pumped into a spherical weather balloon. At any time  $t$ , the volume of the balloon is  $V(t)$  and its radius is  $r(t)$ .  
 (a) What do the derivatives  $dV/dr$  and  $dV/dt$  represent?  
 (b) Express  $dV/dt$  in terms of  $dr/dt$ .

-  **87.** The flash unit on a camera operates by storing charge on a capacitor and releasing it suddenly when the flash is set off.

The following data describe the charge  $Q$  remaining on the capacitor (measured in microcoulombs,  $\mu\text{C}$ ) at time  $t$  (measured in seconds).


$t$	0.00	0.02	0.04	0.06	0.08	0.10
$Q$	100.00	81.87	67.03	54.88	44.93	36.76

- (a) Use a graphing calculator or computer to find an exponential model for the charge.  
 (b) The derivative  $Q'(t)$  represents the electric current (measured in microamperes,  $\mu\text{A}$ ) flowing from the capacitor to the flash bulb. Use part (a) to estimate the current when  $t = 0.04$  s. Compare with the result of Example 2 in Section 2.1.


-  **88.** The table gives the US population from 1790 to 1860.

Year	Population	Year	Population
1790	3,929,000	1830	12,861,000
1800	5,308,000	1840	17,063,000
1810	7,240,000	1850	23,192,000
1820	9,639,000	1860	31,443,000

- (a) Use a graphing calculator or computer to fit an exponential function to the data. Graph the data points and the exponential model. How good is the fit?  
 (b) Estimate the rates of population growth in 1800 and 1850 by averaging slopes of secant lines.  
 (c) Use the exponential model in part (a) to estimate the rates of growth in 1800 and 1850. Compare these estimates with the ones in part (b).  
 (d) Use the exponential model to predict the population in 1870. Compare with the actual population of 38,558,000. Can you explain the discrepancy?

-  **89.** Computer algebra systems have commands that differentiate functions, but the form of the answer may not be convenient and so further commands may be necessary to simplify the answer.

- (a) Use a CAS to find the derivative in Example 5 and compare with the answer in that example. Then use the simplify command and compare again.  
 (b) Use a CAS to find the derivative in Example 6. What happens if you use the simplify command? What happens if you use the factor command? Which form of the answer would be best for locating horizontal tangents?

-  **90.** (a) Use a CAS to differentiate the function

$$f(x) = \sqrt{\frac{x^4 - x + 1}{x^4 + x + 1}}$$

- and to simplify the result.  
 (b) Where does the graph of  $f$  have horizontal tangents?  
 (c) Graph  $f$  and  $f'$  on the same screen. Are the graphs consistent with your answer to part (b)?

91. Use the Chain Rule to prove the following.  
 (a) The derivative of an even function is an odd function.  
 (b) The derivative of an odd function is an even function.

92. Use the Chain Rule and the Product Rule to give an alternative proof of the Quotient Rule.  
 [Hint: Write  $f(x)/g(x) = f(x)[g(x)]^{-1}$ .]

93. (a) If  $n$  is a positive integer, prove that

$$\frac{d}{dx} (\sin^n x \cos nx) = n \sin^{n-1} x \cos(n+1)x$$

- (b) Find a formula for the derivative of  $y = \cos^n x \cos nx$  that is similar to the one in part (a).

94. Suppose  $y = f(x)$  is a curve that always lies above the  $x$ -axis and never has a horizontal tangent, where  $f$  is differentiable everywhere. For what value of  $y$  is the rate of change of  $y^5$  with respect to  $x$  eighty times the rate of change of  $y$  with respect to  $x$ ?

95. Use the Chain Rule to show that if  $\theta$  is measured in degrees, then

$$\frac{d}{d\theta} (\sin \theta) = \frac{\pi}{180} \cos \theta$$

(This gives one reason for the convention that radian measure is always used when dealing with trigonometric functions in calculus: The differentiation formulas would not be as simple if we used degree measure.)

96. (a) Write  $|x| = \sqrt{x^2}$  and use the Chain Rule to show that

$$\frac{d}{dx} |x| = \frac{x}{|x|}$$

- (b) If  $f(x) = |\sin x|$ , find  $f'(x)$  and sketch the graphs of  $f$  and  $f'$ . Where is  $f$  not differentiable?

- (c) If  $g(x) = \sin |x|$ , find  $g'(x)$  and sketch the graphs of  $g$  and  $g'$ . Where is  $g$  not differentiable?


97. If  $y = f(u)$  and  $u = g(x)$ , where  $f$  and  $g$  are twice differentiable functions, show that

$$\frac{d^2 y}{dx^2} = \frac{d^2 y}{du^2} \left( \frac{du}{dx} \right)^2 + \frac{dy}{du} \frac{d^2 u}{dx^2}$$

98. If  $y = f(u)$  and  $u = g(x)$ , where  $f$  and  $g$  possess third derivatives, find a formula for  $d^3 y/dx^3$  similar to the one given in Exercise 97.

## APPLIED PROJECT

### WHERE SHOULD A PILOT START DESCENT?



An approach path for an aircraft landing is shown in the figure and satisfies the following conditions:

- (i) The cruising altitude is  $h$  when descent starts at a horizontal distance  $\ell$  from touch-down at the origin.
- (ii) The pilot must maintain a constant horizontal speed  $v$  throughout descent.
- (iii) The absolute value of the vertical acceleration should not exceed a constant  $k$  (which is much less than the acceleration due to gravity).


1. Find a cubic polynomial  $P(x) = ax^3 + bx^2 + cx + d$  that satisfies condition (i) by imposing suitable conditions on  $P(x)$  and  $P'(x)$  at the start of descent and at touchdown.

2. Use conditions (ii) and (iii) to show that

$$\frac{6hv^2}{\ell^2} \leq k$$

3. Suppose that an airline decides not to allow vertical acceleration of a plane to exceed  $k = 860 \text{ mi/h}^2$ . If the cruising altitude of a plane is 35,000 ft and the speed is 300 mi/h, how far away from the airport should the pilot start descent?

4. Graph the approach path if the conditions stated in Problem 3 are satisfied.

 Graphing calculator or computer required